

Table 1

Anthelmintics available for dogs in France

Tradename	Active	Dose-rate/ regimen	Formulation	Minimum age or weight of treatment	Spectrum of activity (gastrointestinal helminths)		Heartworm preventatives
					Nematocidal	Cestocidal	
Various	salts of piperazine	*	Syrup	Minimum bodyweight : see individual products	Induces the elimination of ascarids (and not their lysis)		
Strongid® chien	Pyrantel	5mg/kg	Paste	2 weeks	Y		
Stronghold®	Selamectin	6mg/kg/ month	Spot on	6 weeks	Y		Y
Advocate®	Moxidectine + Imidacloprid	2.5mg/kg moxidectine + 10mg/kg imidacloprid	Spot on	7 weeks of age or 1 kg	Y		Y
Panacur®	Fenbendazole	50mg/kg*	Tablet	2 weeks	Y	Ta	
Flubenol® Easy Flubenol® Pâte	Flubendazole	22mg/kg for 2-3 days*	Chewable tablet, paste	2 weeks	Y	Ta	
Telmin® KH Telkan®	Mebendazole	50mg/kg for 2-5 days*	Tablet	2 weeks	Y		
Dolthène®	Oxfendazole	11.3mg/kg for 3 days*	Solution	2 weeks	Y	Ta Di	
Vitaminthe®, Polyverkan®	Oxibendazole + Niclosamide	1.5mg/kg oxibend. + 12mg/kg niclosamide	Paste	2 weeks	Y	Ta Di	
Lopatul®, Scanil®, Troscan®	Nitroscanate	50mg/kg	Tablet	2 weeks	Y	Ta Di	
Ascatène®	Pyrantel + Niclosamide	1 tablet per 2-4kg	Tablet		Y	Ta Di	
Stromiten®,	Levamisole +	*	Tablet, Paste	3 months	Y	Ta Di	

Anthelmintic Tables F

Gelminthe®	Niclosamide						
Drontal® P	Pyrantel + febantel + praziquantel	1 tablet per 10 kg	Tablet	2 weeks	Y	Ta Di E	
Interceptor®	Milbemycin oxime	0.5mg/kg	Tablet		Y		Y
Milbemax®	Milbemycin + praziquantel	0.5mg/kg milbemycin oxime + 5 mg/kg praziquantel	Tablet	6 weeks of age or 0.5 kg	Y	Ta Di E	Y
Dolpac®	Pyrantel+ oxantel+ praziquantel	5mg/kg pyrantel+20 mg/kg oxantel+5 mg/kg praziquantel	Tablet	8 weeks of age or 1 kg	Y	Ta Di E	
Droncit®	Praziquantel	5mg/kg	Injection Tablet	Not to unweaned puppies		Y Y	

Y: yes; Ta: *Taenia* spp. E: *Echinococcus* spp; Di: *Dipylidium caninum*

*: see data sheet for more details of dosing regimens

Table 2

Anthelmintics available for cats in France

Tradename	Active	Dose-rate/ regimen	Formulation	Minimum age or weight of treatment	Spectrum of activity (gastrointestinal helminths)		Heart worm preventa tives
					Nematocidal	Cestocida I	
Various	salts of piperazine	*	Syrup	Minimum bodyweight : see individual products	Induces the elimination of ascarids (and not their lysis)		
Strongid® chat, Sépantel® Félin	Pyrantel	5mg/kg	Paste	2 weeks	Y		
Stronghold®	Selamectin	6mg/kg/ month	Spot on	6 weeks	Y		Y
Advocate®	Moxidectine + Imidacloprid	2.5mg/kg moxidectine + 10 mg/kg imidacloprid	Spot on	9 weeks of age or 1 kg	Y		Y
Profender	Emodepside + Praziquantel	3mg/kg emodepside and 12 mg/kg praziquantel	Spot on	8 weeks or less than 0.5 kg	Y	Y	
FlubénoI® Pâte	Flubendazole	22mg/kg for 2-3 days*	Paste	2 weeks	Y	Ta	
Telmin® KH Telkan®	Mebendazole	50mg/kg for 2-5 days*	Tablet	2 weeks	Y		
Vitaminthe®, Polyverkan®	Oxibendazole + Niclosamide	1.5mg/kg oxibend. + 12mg/kg niclosamide	Paste	2 weeks	Y	Ta Di	
Ascatène®	Pyrantel + Niclosamide	1 tablet per 2-4 kg	Tablet		Y	Ta Di	
Stromiten®,	Levamisole +	*	Tablet, Paste	3 months	Y	Ta Di	

Anthelmintic Tables F

Gelminthe®	Nicosamide						
Drontal® chat	Pyrantel + praziquantel	1 tablet per 10 kg	Tablet	2 weeks	Y	Ta Di E	
Milbemax®	Milbemycin + praziquantel	0.5mg/kg milbemycin oxime + 5 mg/kg praziquantel	Tablet	6 weeks of age or 0.5 kg	Y	Ta Di E	Y
Droncit®	Praziquantel	5mg/kg	Injection	Not to unweaned kittens		Y	
			Tablet			Y	

Y: yes; Ta: *Taenia* spp. ; Di: *Dipylidium caninum*

*: See data sheet for more details of dosing regimens