

## Anthelmintics available for dogs in France (May 2009)

Tradename	Active	Dose-rate/ regimen	Formulation	Minimum age or weight of treatment	Spectrum of activity (gastrointestinal helminths)		Heartworm preventatives
					Nematocidal	Cestocidal	
• <b>Advocate®</b>	moxidectine + imidacloprid	2.5 mg/kg moxidectine + 10 mg/kg imidacloprid	Spot on	7 weeks of age or 1 kg	Y		Y
• <b>Ascatène®</b>	pyrantel + niclosamide	1 tablet per 2-4 kg	Tablet		Y	Ta, Di	
• <b>Dolpac®</b>	pyrantel + oxantel + praziquantel	5 mg/kg pyrantel + 20 mg/kg oxantel + 5 mg/kg praziquantel	Tablet	8 weeks of age or 1 kg	Y	Ta, Di, E	
• <b>Dolthène®</b>	oxfendazole	11.3 mg/kg for 3 days*	Solution	2 weeks	Y	Ta, Di	
• <b>Droncit®</b>	praziquantel	5 mg/kg	Injection or tablet	Not to unweaned puppies		Ta, Di, E	
• <b>Drontal® P, XL</b> • <b>Dronstop® chiot</b>	pyrantel + febantel + praziquantel	1 tablet per 10 kg (P) or 35 kg (XP)	Tablet, Solution	2 weeks	Y	Ta, Di, E	
• <b>Flubenol® easy</b> • <b>Flubenol® Pâte</b>	flubendazole	22 mg/kg for 2-3 days*	Chewable tablet, Paste	2 weeks	Y	Ta	
• <b>Interceptor®</b>	milbemycin oxime	0.5 mg/kg	Tablet		Y		Y
• <b>Lopatul®</b>	nitroscanate	50 mg/kg	Tablet	2 weeks	Y	Ta, Di	
• <b>Milbémax® comprimés chien</b>	milbemycin oxime + praziquantel	0.5 mg/kg milbemycin oxime + 5 mg/kg praziquantel	Tablet	6 weeks of age or 0.5 kg	Y	Ta, Di, E	Y
• <b>Panacur® 250 chien</b>	fenbendazole	50 mg/kg*	Tablet	2 weeks	Y	Ta	
• <b>Profender® chien</b>	emodepside + praziquantel	1 mg/kg emodepside + 5 mg/kg praziquantel	Tablet	12 weeks of age	Y	Ta, Di, E	

• <b>Stronghold®</b>	selamectin	6 mg/kg/ month	Spot on	6 weeks	Y		Y
• <b>Stromiten®</b> , • <b>Gelminthe®</b>	levamisole + niclosamide	*	Tablet, Paste	3 months	Y	Ta, Di	
• <b>Strongid® chien</b>	pyrantel	5 mg/kg	Paste	2 weeks	Y		
• <b>Telmin® KH</b>	mebendazole	50 mg/kg for 2-5 days*	Tablet	2 weeks	Y		
• <b>Vitaminthe®</b>	oxibendazole + niclosamide	1.5 mg/kg oxibend. + 12 mg/kg niclosamide	Paste	2 weeks	Y	Ta, Di	
<b>Various names</b>	salts of piperazine	*	Syrup	*	induces the elimination of ascarids (and not their lysis)		

Y: yes; Ta: *Taenia* spp. E: *Echinococcus* spp; Di: *Dipylidium caninum*

\*: see data sheet for more details of dosing regimens

## Anthelmintics available for cats in France (May 2009)

Tradename	Active	Dose-rate/ regimen	Formulation	Minimum age or weight of treatment	Spectrum of activity (gastrointestinal helminths)		Heart worm preventatives
					Nematocidal	Cestocidal	
• <b>Advocate®</b>	moxidectine + imidacloprid	2.5 mg/kg moxidectine + 10 mg/kg imidacloprid	Spot on	9 weeks of age or 1 kg	Y		Y
• <b>Ascatène®</b>	pyrantel + niclosamide	1 tablet per 2-4 kg	Tablet		Y	Ta, Di	
• <b>Droncit®</b>	praziquantel	5 mg/kg	Injection Tablet	Not to unweaned kittens		Ta, Di, E	
• <b>Drontal® chat</b>	pyrantel + praziquantel	1 tablet per 10 kg	Tablet	2 weeks	Y	Ta, Di, E	
• <b>Flubenol® Pâte</b>	flubendazole	22 mg/kg for 2-3 days*	Paste	2 weeks	Y	Ta	
• <b>Milbémax® comprimé chat</b>	milbemycin oxime + praziquantel	0.5 mg/kg milbemycin oxime + 5 mg/kg praziquantel	Tablet	6 weeks of age or 0.5 kg	Y	Ta, Di, E	Y
• <b>Profender® spot on chat</b>	emodepside + praziquantel	3 mg/kg emodepside and 12 mg/kg praziquantel	Spot on	8 weeks or less than 0.5 kg	Y	Ta, Di, E	
• <b>Stromiten®, Gelminthe®</b>	levamisole + niclosamide	*	Tablet, Paste	3 months	Y	Ta, Di	
• <b>Stronghold®</b>	selamectin	6 mg/kg/ month	Spot on	6 weeks	Y		Y
• <b>Strongid® chat</b>	pyrantel	5 mg/kg	Paste	2 weeks	Y		
• <b>Telmin® KH</b>	mebendazole	50 mg/kg for 2-5 days*	Tablet	2 weeks	Y		
• <b>Vitaminthe®</b>	oxibendazole + niclosamide	1.5 mg/kg oxibend. + 12 mg/kg niclosamide	Paste	2 weeks	Y	Ta, Di	
<b>Various names</b>	salts of piperazine	*	Syrup	*	induces the elimination of ascarids (and not their lysis)		

Y: yes; Ta: *Taenia* spp. E: *Echinococcus* spp; Di: *Dipylidium caninum*

\*: See data sheet for more details of dosing regimens